

The role of N-use efficiency in lowering nitrous oxide emissions

James Humphreys¹, Dejun Li^{1,2}, Mingjia Yan², Stan Lalor³, Bernard Hyde⁴,
Gary Lanigan³, Nick Holden², Catherine Watson⁵

¹Teagasc Research and Innovation Centre, Moorepark, Fermoy, Co. Cork.

²School of Agriculture, Food Science and Veterinary Medicine, University College Dublin.

³Teagasc, Johnstown Castle, Wexford.


⁴Environmental Protection Agency, Monaghan.

⁵Agriculture, Food and Environmental Science Division, Agri-Food and Biosciences Institute, Belfast.




Introduction and overview

- Sources of N₂O from Irish agriculture
- Grassland 90% of agricultural area - scale of solution
- The process of denitrification and N₂O losses
- N-use efficiency
- Options
 - nitrification inhibitors
 - slurry application with trailing shoe, injection
 - White clover
- Conclusions








Sources of nitrous oxide emissions from agriculture 2008

(McGettigan *et al.*, 2010)



Nitrification and Denitrification






Soil is an important source of N for plant uptake

Sources of nitrous oxide emissions from agriculture 2008



(McGettigan *et al.*, 2010)



National fertilizer N use




Fertilizer N use on dairy farms (kg/ha)




NFS Dairy Farms		2004	2008
Stocking density (LU per ha)		1.90	1.88
Fertilizer N use (kg per ha)		175	142
Relative fertilizer N use ▼			0.81
Fertilizer N eff. (kg per LU)		92.1	75.5


Cost of fertilizer N (€ per kg) 1990 - 2009



Fertilizer N:milk price ratio 1990 - 2009



Fertilizer N costs and national use



National fertilizer N use

- Closely related to fertilizer N cost
- Projected ▲ in fertiliser N cost
- Especially relative to product prices
- Anticipate ▼ in national fertilizer N use
- Associated ▼ in N₂O emissions
- Alternatives to fertilizer N


Making more efficient use of fertilizer N on farms

- Farm gate N balance: surplus of N in inputs minus outputs in products
- Typical range in N-use efficiency = 15 to 35%



- Increase the efficiency of recycling within the farm gate (Treacy *et al.*, 2008)
- Avoid wasteful use imported fertilizers – wrong times, matching rates to req. etc.
- ▼ farm gate surplus by 1 kg N per t milk = 29 kg CO₂ eq. (Schils *et al.* 2006)

Nitrification and Denitrification: Nitrification inhibitors




Inhibiting nitrification can lower N_2O emissions (and nitrate leaching)

Table 8: Emissions of nitrous oxide after application of different nitrification inhibitors with different fertilisers.


Nitrification inhibitor	Fertiliser	Country	Crop	Reduction of N ₂ O losses as %	Length of monitoring	Author
DMPP	Liquid cattle slurry	Spain	Clover/ryegrass	29	59 days	Menendez <i>et al.</i> (2006)
DMPP	Liquid cattle slurry	Spain	Pasture	69	106 days (autumn)	Merino <i>et al.</i> (2005)
				48	29 days (spring)	
DMPP	Ammonium sulphate nitrate	Spain	Grass	58	20 days	Macadam <i>et al.</i> (2003)
	Liquid cattle slurry			61	60 days	
DMPP	Liquid cattle manure	UK	Grass	32	22 days	Dittert <i>et al.</i> (2001)
DMPP	Ammonium sulphate nitrate	Germany	Winter wheat	20	4 months	Linzmeier <i>et al.</i> (2001)
DMPP	Ammonium sulphate nitrate	Germany	Barley	41	3 years	Weiske <i>et al.</i> (2001)
			Maize	47		
			Wheat	53		
DCD			Barley	30	3 years	
			Maize	22		
			Wheat	29		
DCD	Urine	New Zealand	Grass	52	2-3 months (autumn)	Zaman <i>et al.</i> (2009)
				39	2-3 months (spring)	
				16.5	2-3 months (summer)	
DCD	Urea	New Zealand	Grass	75-91	2-3 months	Smith <i>et al.</i> (2008)
DCD	Urine	New Zealand	Grass	61-73	69-137 days	Di <i>et al.</i> (2007)
DCD	Urine	New Zealand	Grass	82	16 months	Di and Cameron, (2002)
DCD	Ammonium sulphate	India	Rice	26	98 days	Kumar <i>et al.</i> (2000)
	Urea			11		
DCD	Urea	USA	Spring barley	82	90 days	Delgado & Mosier, 1996
DCD	Urine	Netherlands	Grass	50-89	14 days	DeKlein and van Logtestijn, (1994)
DCD	Ammonium sulphate	UK	grass	40	64 days	Skiba <i>et al.</i> (1993)
Nitrapyrin	Urea	USA	Corn	40-60	100 days	Bronson <i>et al.</i> 1992

DCD and N₂O emissions (RSF funded projects)

Dennis et al., 2008; Teagasc, Johnstown Castle



RSF07-519: Watson et al. unpub., AFBI, UCD, Teagasc



Nitrification inhibitors

- Nitrification inhibitors substantially lower emissions from fertilizer, slurry & urine
- Little or no improvement in agronomic performance (e.g. O'Conner et al., 2010)
- Costs associated with DCD exceed any economic benefit (Monaghan et al., 2009)
- No incentive for Irish farmers to use nitrification inhibitors at present
- Evidence of higher emissions from CAN compared with urea under damp conditions



Slurry application techniques



- Trailing shoe increases N-use efficiency and herbage production etc. compared with splash-plate (Lalor, 2008)
- Trailing shoe increases N_2O emissions because more N enters soil (less NH_3 lost)
- Need more fertilizer N with splash-plate for same level of productivity
- On balance – no benefit or detriment with trailing shoe for N_2O emissions (Huijsmans and Schils, 2009)








White clover



Biological N fixation

Biological N Fixation (BNF) is a photosynthetically derived process where atmospheric N is reduced to ammonium (plant-available form of N) by N-fixing bacteria







White clover and biologically fixed N (BFN)




- Direct N₂O emissions from white clover not included in current IPCC methodology
- Corré and Kasper (2002) BNF emission factor = 0.2% versus 1.3% for fertilizer N
- Conclusion of recent review for IPCC: No direct N₂O emissions should be attributed to BNF (de Klein *et al.*, 2006)
- Photosynthetically derived BNF is “greenhouse gas neutral” (Ledgard *et al.*, 2009)
- Upstream CO₂ emissions:
 - fertilizer N manufacture requires 60 MJ/kg N (Jenssen and Konhshaug, 2003)
 - manufacture, transport and spreading = 3.5 to 4.0 kg CO₂ eq. per kg Urea-N (Williams *et al.*, 2006)
- Need LCA to fully determine differences between systems


1 t N needs 10 barrels of oil




N₂O-N emissions from grassland in New Zealand (Ledgard et al., 1999)



N₂O emissions from grassland in Australia (Eckard et al., 2003)



N₂O emissions from clover-based and N-fertilized grassland at Solohead (Li et al., unpublished, RSF 07 516)




Life Cycle Assessment (LCA) methodology





Primary resources

Raw materials
Energy use
Transport




Inputs

Fertilizers
Electricity
Fuel
Feeds




GHG emissions from Irish clover versus fertilizer N dairy systems

(Humphreys et al. 2009; estimated using LCA methodology; Yan et al. RSF07 516)




Profitability of clover and fertilizer N-based dairy production

(Data from Humphreys *et al.*, 2008; 2009; Mihailescu *et al.* Interreg IVB Dairyman)




Fertilizer N for clover
90 kg per ha




Profitability of clover and fertilizer N-based dairy production

(Data from Humphreys *et al.*, 2008; 2009; Mihailescu *et al.* Interreg IVB Dairyman)



Implications

- Same net margin with ▼ 0.3 LU per ha and with ▼ fertilizer N
- ▼LU = ▼ methane and ▼ nitrous oxide (N excretion during grazing)
- ▼ fertilizer N = ▼ direct and indirect N₂O emissions




GHG emissions from Dutch clover versus fertilizer N dairy systems

(Schils et al., 2000; Schils et al., 2005)

Two systems: 34.4ha and 40.6 ha; 59 cows per system + followers


kg CO₂ eq. per kg energy corrected milk (ECM)

per kg milk = ▼12.5%




GHG emissions from Dutch clover versus fertilizer N dairy systems

(Schils et al., 2000; Schils et al., 2005)




Summary and Conclusions

- Lower N₂O emissions due to less livestock and more efficient N use
- Pressure to increase N use efficiency on farms
- Benefits from current best management practices
- Little incentive to use nitrification inhibitors
- Trailing shoe offer no specific advantages for N₂O emissions
- Clover-based systems offer substantial potential to lower N₂O emissions



“How do we increase food production while reducing emissions”

- Lower N₂O emissions due to less livestock and more efficient N use
- Pressure to increase N use efficiency on farms
- **Benefits from current best management practices**
- Little incentive to use nitrification inhibitors
- Trailing shoe offer no specific advantages for N₂O emissions
- **Clover-based systems offer substantial potential to lower N₂O emissions**



“How do we increase food production while reducing emissions”

- Potential for further cuts in emissions
- Rising cost of energy and fertilizer
- Projecting forward fertilizer N use = 4.5% ▼ in emissions from agriculture
≈ 1% ▼ in national emissions
- Important to get the message out to farmers (Teagasc)
- Schemes – REPS-4 and AEOS
- Funding, expertise and recruitment embargos

