

Curtins Farm Walk Notes Tuesday 02-03-10

Low Stocking Rate Group (2.5 HF Cows/ha)

Critical Issues

- 1) Maintain post-grazing height at between 4 – 5cm
- 2) Avoid Poaching
- 3) Achieve high grass intakes

Situation

Figure 1. Low Stocking Rate Feed Wedge

Teagasc, Dairy Production Department, Moorepark		GrazePlan - Grass Measurement Report	
Group : TEAGASC RESEARCH FARMS		Date Produced 03-MAR-10	
Client : CRT - Aidan Brennan		Effect of stocking rate and calving date on animal performance	
Date : 01-MAR-10		Treatment : Low SR	
Rotation Length :	55	Farm Cover (kg DM/ha) :	518
Grass Allocation /cow (kg grass dry matter/LU)	12	Farm Cover (kg DM/LU) :	206
Concentrate Fed (kg/cow) :	6	Current Monthly Fertilizer Rate (kg/ha) :	
Silage Fed (kg DM/cow) :	0	Stock Rate (LU/ha) :	2.52
N Application Rate (units/acre) :		Growth Rate :	30
N Application Rate (kg/ha) :		Farm Demand (kg DM/LU/day) ::	80
Residual Height :	3.5	Target pregrazing yield (kg DM/ha) :	1661
Total Livestock :	23		





Table 1. Revised Spring Feed Budget for Low Stocking Rate

Week	Grass Cover (kg dm/ha)	Cover per LU	Growth (kg dm/day)	No. of Cows
09/02/09	647	259	6	10
16/02/09	636	254	12	15
23/02/09	626	250	14	18
02/03/09	599	240	15	21
09/03/09	545	218	17	21
16/03/09	486	194	21	21
23/03/09	438	175	23	22
30/03/09	378	151	27	22

Figure 2. Spring Rotation Plan for Low Stocking Rate



- 1) 78% of the herd is calved. Grazing commenced on the 31st of January and 34% of the farmlet was grazed on the 1st of March.
- 2) Farm cover is 518kg/ha (Figure 1) which is below target based on the revised spring budget, (the budget was revised to take into account the lower than expected opening farm cover and lower growth rates due to the cold weather).
- 3) Grass allocation is based on the Spring Rotation Plan (Figure 2). This group is being allocated 0.17ha/day.
- 4) Cows are currently grazing block 15, pre-grazing yield is 1200kg and 0kg residual is left after grazing. Therefore, grass intake is estimated to be 11.3kg/day, [(1200kg allowance*0.17ha)/18 cows calved]. Concentrate supplementation is at 6kg/day.

High Stocking Rate Group (3.3 HF Cows/ha)

Critical Issues

1. Maintain post-grazing height at between 3 and 3.5cm
2. Avoid poaching
3. Achieve high grass intakes

Situation

Figure 3. High Stocking Rate Feed Wedge

Teagasc, Dairy Production Department, Moorepark		GrazePlan - Grass Measurement Report	
Group : TEAGASC RESEARCH FARMS		Date Produced 03-MAR-10	
Client : CRT - Aidan Brennan		Effect of stocking rate and calving date on animal performance	
Date : 01-MAR-10		Treatment : High SR	
Rotation Length :	60	Farm Cover (kg DM/ha) :	484
Grass Allocation /cow (kg grass dry matter/LU)	7.5	Farm Cover (kg DM/LU) :	147
Concentrate Fed (kg/cow) :	6	Current Monthly Fertilizer Rate (kg/ha) :	
Silage Fed (kg DM/cow) :	4	Stock Rate (LU/ha) :	3.30
N Application Rate (units/acre) :		Growth Rate :	11
N Application Rate (kg/ha) :		Farm Demand (kg DM/LU/day) ::	95
Residual Height :	2.8	Target pregrazing yield (kg DM/ha) :	1483
Total Livestock :	23		





Figure 4. Revised Spring Feed Budget for High Stocking Rate

Week	Grass Cover (kg dm/ha)	Cover per LU	Growth (kg dm/day)	No. of Cows
09/02/09	610	186	6.0	10
16/02/09	602	184	13.0	15
23/02/09	603	184	15.4	18
02/03/09	585	178	16.8	21
09/03/09	535	163	18.7	21
16/03/09	477	145	23.0	21
23/03/09	428	131	25.0	22
30/03/09	361	110	29.5	22

Figure 5. Spring Rotation Plan for High Stocking Rate



- 70% of the herd is calved and grazing commenced on the 10th of February, 10 days later than planned. 27% of the farmlet was grazed on the 1st of March.
- Pre-grazing yield is 1170kg and area allocation is 0.11ha/day. Therefore, grass intake is estimated to be 8kg [(1170kg*0.11ha)/16 cows calved]. 6kg of concentrate is being fed along with 4kg of high quality bale silage.
- Farm cover is 484kg/ha which is well behind our original and revised target.
- Cows are on 24-hour allocation of grass. They get fresh grass after evening milking and go back to the same section after morning milking and are then brought indoors for silage once the desired residual is reached, usually around 11am.

Whole Farm Situation

1. Average soil temperature for the past week was 2.9°C.
2. Total rainfall for the week was 5.6mm.
3. Average weekly growth this week was around 11kg/day, average for the previous 3 years was 21kg/day.
4. Dry matters were 27% on Monday morning.
5. 23 units urea/acre was spread on 70% of the farm on 1/02/10 while 2500gls/acre of watery slurry was spread on the remaining 30%.
6. 23units of urea/acre will be spread towards the end of the week on the 705 of the farm that has not been grazed. The grazed paddocks have all received slurry.
7. Latest milk quality test results from the milk processor are; Fat 4.21%, Protein 3.32%, Lactose 4.74%, SCC 171k.
8. Freshly calved cows are on 4kg of concentrate for 5 days after calving, this is then increased to 6kg thereafter.
9. 1 cow was treated for suspected grass tetany and 1 cow was treated for mastitis this week.
10. As we are expecting a grass deficit in the coming weeks we are trying to preserve silage stocks. For this reason all dry cows are now on a diet of 6kg soya hulls and 4kg of wheaten straw.
11. Critical Short-term Actions:
 - a. Cows getting silage are on 24-hour allocations while cows not getting silage are on 12-hour allocations.
 - b. Avoid poaching, we are not back fencing at the moment but if weather deteriorates we will.
 - c. Monitor cows closely for signs of ill health and treat early.