

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Amistar Opti (PCS No. 05068)

CROPS: [Outdoor grown: Asparagus]

This approval provides for the use of the product **Amistar Opti (PCS No. 05068)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 10.05.18.

[Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Amistar Opti
<i>Active Substance:</i>	Azoxystrobin & Chlorothalonil
<i>Formulation:</i>	80g/ l & 400g/ l SC
<i>Authorisation Holder:</i>	Syngenta UK ltd
<i>Marketing Company:</i>	Syngenta Ireland Ltd
<i>PCS No:</i>	05068
<i>User type/Function:</i>	Professional Use Only / Fungicide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Outdoor asparagus	2.5 l/ha	n/a	2.5 l/ha	8 to 12 days	See specific restriction No. 8	Rust and stemphylium

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed sprayer and hand held sprayer only.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (6) This approval extends to professional use only.
- (7) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (8) Application may only be made after the final harvest & before senescence in the year of application. Application may also be made in establishment crops 3 weeks after planting the crowns providing no crop is harvested that year.
- (9) When this application is applied via hand held sprayer, the spray solution must not be more dilute than 420 ml of product in 100 litres of water.
- (10) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Advisory information:

This approval relates to use of Amistar Opti as a horticultural fungicide for use on the crops & situations listed above. Amistar Opti will also give moderate control of botrytis.

Resistance Management:

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, FRAC group] with satisfactory efficacy against the target.

Use Amistar Optic as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action (FRAC group)

To avoid the likelihood of resistance developing, application of Azoxystrobin should be made with due regard to the use of QoI compounds.

Amistar Opti should be used preventatively and should not be relied upon for its curative potential.

Amistar Opti is a co-formulated mixture of the strobilurin azoxystrobin and chlorothalonil. The two actives possess different modes of action and thus form part of a resistance management strategy.

Azoxystrobin is classed FRAC group 11

Chlorothalonil is classed FRAC group M5

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.