

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Decis Protech (PCS No. 05269)

CROPS: Outdoor crops of: Fodder beet, beetroot, carrots, celeriac, horseradish, leeks, salad onion, parsley root, parsnip, radish, rye, durum wheat, salsify, triticale, cress, endives, lambs lettuce, baby leaf, herbs, blackberry, grass (seed), strawberry, currants (white, red & black), gooseberry, potato (seed), bulb onion, garlic, shallot, sweetcorn, chinese cabbage, kohlrabi, linseed, broccoli.

Protected crops of:, celery, rhubarb, strawberry, tomato, aubergine, cucumber, courgette, gherkin, squash, lettuce, lambs lettuce, baby leaf, cress, herbs, mushroom

This approval provides for the use of the product **Decis Protech (PCS No. 05269)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor uses.

The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 14.01.19. [Includes latest changes. This version should be used until updated]

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Product name: Decis Protech

Active Substance: Deltamethrin

Formulation: 15g/l EW

Authorisation Holder: Bayer CropScience Ltd.

Marketing Company: Bayer CropScience Ltd.

PCS No: 05269

User type/Function: Professional Use Only / Insecticide.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Durum wheat, rye and triticale	420 ml/ha	2 per year	840 ml/ha	14 days	30 days before harvest	Aphids, cereal flea beetle, yellow cereal fly, gout fly, saddle gall midge
Grassland (seed crops)	420 ml/ha	3 per year	1.26 l/ha	-	-	Aphids
Fodder beet	500 ml/ha	-	500 ml/ha	-	30 days before harvest	Flea beetle
Outdoor leek	420 ml/ha	-	1.26 l/ha	14 days	3 days before harvest	Cutworm, leaf moth
Outdoor & protected salad onion	420 ml/ha	3 per year	1.26 l/ha	14 days	3 days before harvest	Leaf miner

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Carrot, parsnip, horseradish, celeriac, beetroot, parsley root, radish, salsify	500 ml/ha	3 per crop	1.5 l/ha	14 days	21 days before harvest	Carrot fly, cutworm, thrips
Outdoor crops of Cress, endive, Lambs lettuce, Land cress, baby leaf, herbs	420 ml/ha	3 per crop (6 per year in multiple cycles)	1.26 l/ha	14 days	7 days before harvest	Aphids, caterpillar, flea beetle
Outdoor blackberry	830 ml/ha	3 per year	2.49 l/ha	14 days	7 days before harvest	caterpillar, raspberry beetle
Outdoor strawberry	500 ml/ha	2 per crop	1.0 l/ha	14 days	3 days before harvest	Blossom weevil, caterpillar, Thrips
Outdoor currants (white, red & black), gooseberry	400 ml/ha	2 per crop	800 ml/ha	14 days	7 days before harvest	caterpillar, capsids
Seed potatoes	670 ml/ha	2 per crop	1.34 l/ha	-	7 days before harvest	PLRV transmission by aphid vectors, Colorado beetle
Bulb onion, garlic, shallot	500 ml/ha	2 per crop	1.0 l/ha	14 days	7 days before harvest	Leek moth aphids, thrips, leaf miners
Outdoor sweetcorn	830 ml/ha	3 per crop	2.49 l/ha	-	3 days before harvest	Aphids, caterpillar
Outdoor broccoli, Chinese cabbage, Kohl rabi	500 ml/ha	2 per crop	1.0 l/ha	14 days	7 days before harvest	caterpillar, tortrix moth
Linseed	500 ml/ha	1 per crop	500 ml/ha	-	GS 81	Thrips

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Protected strawberry	830 ml/ha	3 per crop	2.49 l/ha	14 days	3 days before harvest	Blossom weevil, caterpillar Thrips (including western flower thrips)
Protected crops of celery, rhubarb	500 ml/ha	3 per crop	1.5 l/ha	-	7 days before harvest	Caterpillar
Protected crops of tomato, aubergine, cucumber, courgette, gherkin, squash	830 ml/ha	3 per crop	2.49 l/ha	7 days	3 days before harvest	Whitefly, Aphids caterpillar, thrips, tortrix moth
Protected lettuce, lambs lettuce, baby leaf, cress, herbs	830 ml/ha	2 per crop (6 per year in multiple cycles)	1.66 l/ha	7 days	7 days before harvest	Aphids, caterpillar
Mushrooms (Via cold fogging equipment only)	5ml /100m ³ air space	2 per crop	-	14 days	3 days before harvest	Phorid flies, Sciarid flies

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application may only be made using a tractor mounted/trailed sprayer, orchard blast air assisted sprayer, knapsack. Cold fogging equipment (for mushrooms only).
- (4) Post application no unprotected person should enter the treated area until the application had dried & the area has been well ventilated.
- (5) This approval extends to professional use only.
- (6) For environmental information refer to the product label but specifically:
When applying by tractor mounted trailed sprayer: To protect aquatic organisms respect an unsprayed buffer zone of 7m to surface water bodies

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

When applying by air-assisted sprayer to outdoor blackberries:

To protect aquatic organisms respect an unsprayed buffer zone of 30m to surface water bodies

When applying by knapsack sprayer: To protect aquatic organisms respect an unsprayed buffer zone of 1m to surface water bodies. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.

(7) For operator protection refer to the product label.

(8) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Decis as an agricultural/horticultural insecticide on the crops & situations listed above.

Resistance Management:

Use Decis as part of an Integrated Crop Management (ICM) strategy.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical groups or alternative control measures should be included in the planned anti-resistance strategy. This product should be used in conjunction with other fungicides from different cross-resistant groups [different mode of action, IRAC group] with satisfactory efficacy against the target.

Decis contains Deltamethrin IRAC class: 3A.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.