

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)

INSTRUCTIONS FOR USE

PPP: Escolta (PCS No. 05681)

CROPS:

[Outdoor Grown: Beetroot]

This approval provides for the use of the product **Escolta (PCS No. 05681)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor use. The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 15.02.18. [Includes latest changes. This version should be used until updated]

| | |
|------------------------------|---------------------------------|
| <i>Product name:</i> | Escolta |
| <i>Active Substance:</i> | Cyproconazole & Trifloxystrobin |
| <i>Formulation:</i> | 160g & 375g/l SC |
| <i>Authorisation Holder:</i> | Bayer CropScience Limited |
| <i>Marketing Company:</i> | Bayer CropScience Limited |
| <i>PCS No:</i> | 05681 |
| <i>User type/Function:</i> | Professional Use / Fungicide. |
| <i>Application:</i> | |

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

| Crops: | Max. individual dose | Max. number of applications | Max. total dose | Minimum interval between applications | Latest time of application | Target |
|---------------------------|----------------------|-----------------------------|-----------------|---------------------------------------|----------------------------|--|
| [Outdoor Grown: beetroot] | 0.35 l/ha | 2 per crop | 0.70 l/ha | 21days | 21 days before harvest | Powdery Mildew, Rust, Cercospora & Ramularia |

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) Application by tractor mounted/trailed/air assisted sprayer only.
- (4) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (5) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water. Growers may be eligible to reduce the buffer zone under the PRCD STRIPE Scheme.
- (6) The first application must not be made before BBCH 31. If a second application is required, it must not be made before BBCH 40
- (7) This approval extends to professional use only.
- (8) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (9) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Escolta as a horticultural fungicide for use on the crops and situations listed above.

Growers are advised to test a small area of the crop first to ensure the application will not cause damage to the crop especially if the crop is under any stress.

Growth stage BBCH31 indicates that 10% of the soil is covered by the crop's leaves. BBCH 39 refers to 90% leaf cover. BBCH 40 indicates the start of root swelling.

Resistance Management:

This product should be applied in accordance with an anti-resistance strategy. Disease control may be reduced if strains of the pathogen less sensitive to this active develop.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical types or alternative control measures should be included in the planned programme. Alternating pesticides with different modes of action is a recognised anti-resistance strategy and this product should always be used in conjunction with other products with a different mode of action / different cross-resistance group with satisfactory efficacy against the target.

Escolta containing Cyproconazole is classed by FRAC as group 3 triazoles. Escolta containing Trifloxystrobin is classed by FRAC as group 11 QoL fungicides.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.
