

EXTENSION OF AUTHORISATION FOR MINOR USES (EAMU)**INSTRUCTIONS FOR USE****PPP: Nativo 75 WG (PCS No. 04642)**

CROPS:

Outdoor crops of parsnip, celeriac, celery, horseradish, parsley root, salsify, leek, salad onion, swede, and turnip.

This approval provides for the use of the product **Nativo 75 WG (PCS No. 04642)** in respect of, the crops and situations as listed above, which are extra to those specified on the product label.

The conditions of application outlined below are statutory. They must be complied with when availing of this extension for minor use. The instructions provided in this extension for minor uses must be used in conjunction with information provided on the approved product label.

Failure to abide by the conditions of approval specified below or on the product label may constitute a breach of the product approval.

Reference Date: 06.11.18. [Includes latest changes. This version should be used until updated]

<i>Product name:</i>	Nativo 75 WG
<i>Active Substance:</i>	Trifloxystrobin 250g/kg Tebuconazole 500g/kg
<i>Formulation:</i>	Water dispersible granule
<i>Authorisation Holder:</i>	Bayer CropScience Limited
<i>Marketing Company:</i>	Bayer CropScience limited
<i>PCS No:</i>	04642
<i>User type/Function:</i>	Professional Use / Fungicide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

Application:

Crop	Max. individual dose	Max. number of applications	Max. total dose	Minimum interval between applications	Latest time of application	Target
Parsnips, parsley root, horseradish, salsify	0.3 kg/ha	3 per crop	0.9 kg/ha	21 days	21 days before harvest	Alternaria, Sclerotinia, powdery mildew
Celery	0.3 kg/ha	3 per crop	0.9 kg/ha	21 days	21 days before harvest	Black blight (<i>Septoria apiicola</i>)
Celeriac	0.3 kg/ha	2 per crop	0.6 kg/ha	21 days	21 days before harvest	(<i>Mycosphaerella Brassicicola</i>), alternaria (<i>Alternia brassicae</i>), Black blight (<i>septoria apiicola</i>), powdery mildew
Leeks, salad onion	0.36 kg/ha	1 per crop	0.36 kg/ha	21 days	21 days before harvest	Rust
Swedes, turnip	0.36 kg/ha	2 per crop	0.72 kg/ha	21 days	21 days before harvest	<i>Mycosphaerella Brassicicola</i>), alternaria (<i>Alternia brassicae</i>), Black blight (<i>septoria apiicola</i>), powdery mildew

Other specific restrictions:

- (1) To avoid risks to human health & the environment comply with the terms of this approval in conjunction with the directions for use on the product label.
- (2) For any individual crop the maximum total dose refers to the total dose of any product containing the same active ingredient.
- (3) A maximum number of applications of three per calendar year on a treated area must not be exceeded
- (4) Application by tractor mounted/trailed sprayer only. This product must not be applied via hand-held equipment

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

- (5) Post application, do not allow unprotected persons to enter treated areas until the crop is completely dry & has been well ventilated.
- (6) For environmental information refer to the product label but specifically: To protect aquatic organisms respect an unsprayed buffer zone of 30m to surface water bodies from tractor mounted/trailed sprayers. Aim spray away from water.
- (7) This approval extends to professional use only.
- (8) For operator protection refer to the product label in conjunction with the specific restrictions above.
- (9) Adverse effects: Approval holders must immediately submit any new information on the potentially dangerous effects of a product or residues of an active substance contained in a product, on human or animal health, or the environment.

Advisory information:

This approval relates to the use of Nativo 75WG as an agricultural fungicide for use on the crops and situations listed above.

Applications should be applied before disease establishes itself in the crop (making use of disease forecasting systems as appropriate) using a minimum of 300 litres of water per ha.

Growers are advised to test a small area of the crop first to ensure the application will not cause damage to the crop especially if the crop is under any stress.

Resistance Management:

This product should be applied in accordance with an anti-resistance strategy. Disease control may be reduced if strains of the pathogen less sensitive to Trifloxystrobin & Tebuconazole develop.

Total reliance on one pesticide will hasten the development of resistance. Pesticides of different chemical types or alternative control measures should be included in the planned programme. Alternating fungicides with different modes of action is a recognised anti-resistance strategy and Nativo 75WG should always be used in conjunction with other fungicides with a different mode of action / different cross resistance group with satisfactory efficacy against the target.

Trifloxystrobin is classed (FRAC 11) & is a QoI fungicide.

Tebuconazole is classed (FRAC 3) & is a DMI fungicide.

There is limited evidence of crop safety and/or product efficacy available for this extension of authorisation for minor uses and the commercial risk of using this product on the crops/situations listed in this document is borne entirely by the grower.

IMPORTANT PLEASE NOTE:

Prior to the use of this product, growers should take advice from their crop advisors particularly in relation to the use of this product within an integrated crop management system of production.
